

Attenuation of High Glucose-Induced Rat Cardiomyocyte Apoptosis by Exendin-4 *via* Intervention of HO-1/Nrf-2 and the PI3K/AKT Signaling Pathway

Shu-Mei Zhao¹, Hong-Li Gao¹, Yong-Liang Wang¹, Qing Xu², and Chun-Yan Guo¹

¹Cardiovascular Center

Beijing Friendship Hospital, Capital Medical University, Beijing 100050

and

²College of Basic Medicine, Capital Medical University, Beijing 100069, People's Republic of China

Abstract

Exendin-4, a glucagon-like peptide-1 receptor agonist, demonstrated cytoprotective actions beyond glycemic control in recent studies. The aims of the present study were to investigate the effects of exendin-4 on high glucose (HG)-induced cardiomyocyte apoptosis and the possible mechanisms. Rat cardiomyocytes were divided into 3 groups: normal glucose group (NG group), HG group and HG +exendin-4 group (HG+Ex Group). Cardiomyocyte apoptosis was evaluated by double-staining with annexin V-fluorescein isothiocyanate (FITC)/propidium iodide (PI) and flow cytometry. Intracellular reactive oxygen species (ROS) production was detected by 2',7'-dichlorodihydrofluorescein diacetate (DCHF-DA) incubation and fluorescence microscopy. LY294002 (LY), a phosphoinositide 3-kinase (PI3K) pathway inhibitor, was added to the medium of the HG+Ex+LY Group for further western blot analysis. The proteins analyzed involved oxidative stress-associated proteins, heme oxygenase-1 (HO-1) and nuclear factor E2-related factor 2 (Nrf-2), and apoptosis-associated proteins, caspase-3, Bax/B-cell lymphoma 2 (Bcl-2) and p-AKT/AKT. HG treatment induced cardiomyocyte apoptosis ($P = 0.00$) and clearly upregulated ROS production ($P = 0.00$); exendin-4 co-incubation also ameliorated cardiomyocyte apoptosis ($P = 0.004$) and decreased ROS ($P = 0.00$) level significantly. HO-1 and Nrf-2 protein expression levels decreased significantly in the HG group ($P < 0.05$), but the levels were elevated by exendin-4 intervention ($P < 0.05$). Furthermore, exendin-4 attenuated HG-induced higher protein expression, including cleaved caspase-3 and Bax, increased the expression of Bcl-2 protein ($P < 0.05$). However, these impacts of exendin-4 were counteracted significantly by co-incubation with LY294002. In addition, exendin-4 ameliorated HG-induced p-AKT/AKT lower expression, and this impact was also suppressed by LY294002. Exendin-4 ameliorates HG-induced cardiomyocyte apoptosis, and the mechanisms may involve anti-oxidative stress *via* the HO-1/Nrf-2 system, as well as intervention of the PI3K/AKT signaling pathway.

Key Words: cardiomyocyte apoptosis, exendin-4, high glucose, HO-1/Nrf-2 system, PI3K/AKT pathway

Introduction

With the high prevalence of diabetes, more attention has been paid to diabetic cardiomyopathy.

Diabetes-associated myocardial impairment was initially recognized by Rubler *et al.* (17) in diabetic patients, who exhibited defects in both electrical and mechanical properties, and increased morbidity

and mortality. Myocardial contraction and relaxation in diabetic models are markedly impaired, and the cardiac dysfunction, and even congestive heart failure, may result from a variety of metabolic and biochemical abnormalities. Exploration of the mechanisms and intervention strategies, involving oxidative stress and alteration in Ca^{2+} homeostasis, has been always the hot topics in this field (16, 29).

Glucagon-like peptide-1 (GLP-1) is an incretin hormone secreted by the small intestine in a nutrient-dependent manner, which stimulates insulin secretion and inhibits glucagon secretion and gastric empty (2, 12). GLP-1 has wide-range effects on glucose metabolism, and the GLP-1 pathway has been a therapeutic target in the treatment of diabetes (4, 10). However, GLP-1 has a short half-life of <2 min in the circulation, and it is degraded rapidly by the enzyme dipeptidyl peptidase-4 (DPP-4), limiting the biological efficacy in clinical practice. Current treatment *via* the GLP-1 pathway focuses on increasing the level of GLP-1 in the blood, including the use of either compounds to inhibit DPP-4 activities, or application of GLP-1 receptor agonists. Exendin-4, a GLP-1 receptor agonist, is a potential insulin secretagogue, and has been approved for the treatment of type 2 diabetes (4, 10). Furthermore, exendin-4 has a therapeutic advantage in its resistance to degradation by DPP-4, resulting in a longer pharmacological half-life (4, 10).

Beyond glycemic control, cytoprotective actions of the GLP-1 pathway involves protection of pancreatic beta cells (23) and neurons (5). A variety of cardiovascular benefits of the GLP-1 pathway has been reported, including protection against post-myocardial infarction remodeling (15), improvement of cardiac function in mice over-expressing monocyte chemoattractant protein-1 in cardiomyocytes (26), and amelioration of cardiac ischemia/reperfusion injury (22). Although these cardioprotective effects *via* the GLP-1 pathway have been reported, the related mechanisms of these effects are still unclear.

It is well known that oxidative stress is responsible for the induction of cardiomyocytes apoptosis *via* high glucose (HG). Mangmool, S. *et al.* (11) demonstrated that stimulation of the GLP-1 receptor with exendin-4 attenuated H_2O_2 -induced reactive oxygen species (ROS) production and increased the synthesis of antioxidant enzymes. Hence, we hypothesized that exendin-4 may protect diabetic myocardium and ameliorate cardiomyocyte apoptosis, and the mechanisms may involve oxidative stress and apoptosis signaling pathway. The aims of the present study were to investigate the effects of exendin-4 on HG-induced cardiomyocyte apoptosis, and the roles of exendin-4 in the intervention of oxidative stress and the PI3K/AKT signal pathways in these pathologic processes.

Materials and Methods

Cell Isolation and Culture

The animal experimental protocols were approved by the Beijing Friendship Hospital Animal Care and Research Committee. All surgery was performed under anesthesia, and all efforts were made to minimize sufferings. Cardiomyocytes were prepared by enzymatic disassociation of 1- to 2-day old Sprague-Dawley rats ($n = 8-10$) as described previously (9). Briefly, the rats were sacrificed by opening the pleura under ether anesthesia and aseptic conditions. The hearts of the rats were removed rapidly and cleaned in iced-cold D-Hanks balanced salt solution. The minced ventricular myocardium was digested with trypsin (0.07%) and collagenase II (0.04%) at 37°C . Cells were extracted by repeated tissue digestion and the whole-cell suspensions were centrifuged at 1,000 rpm for 10 min. After the removal of supernatants, the cell pellets were re-suspended in Dulbecco's modified eagle medium (DMEM), which contained 15% fetal bovine serum and 1% penicillin-streptomycin. The cells were plated onto 10-cm culture dishes at 37°C , 5% CO_2 for 90 min. When the non-myocytes attached to the dishes, the cardiomyocytes remained in the suspension and were harvested and seeded in six-well culture plates coated with laminin. After incubation at 37°C , 5% CO_2 for 24 h, the cardiomyocytes were identified by α -actinin staining, and were divided into 3 groups: normal glucose group (NG Group: DMEM containing glucose 5.5 mM; Hyclone, MA, USA), HG Group (DMEM containing glucose 25 mM), HG + exendin-4 group (HG+Ex Group) (exendin-4 purchased from Sigma, St. Louis, MO, USA). The vehicle for exendin-4 was phosphate buffer saline (PBS). Cardiomyocytes were cultured in serum-free DMEM for 12 h before experiments. Cells in different groups were incubated at 37°C , 5% CO_2 for 24 h before further study assessments.

Cell Survival Analysis

Cell counting kit-8 (CCK-8, DOJINDO, Japan) was used to evaluate the effects of HG and exendin-4 on the cell viability, as required by manufacturer's protocol. Cardiomyocytes were seeded into 96-well dishes, and the density was adjusted to 5×10^4 cells/ml. Cells were incubated in culture media with NG (5.5 mM), HG (25 mM) or HG+Ex at 10 nM, 20 nM, 30 nM or 40 nM at 37°C , 5% CO_2 for 24 h. CCK-8 (10 μl) was then added to each well and incubated for another 4 h. The absorbance at 450 nm was measured using a microplate reader (BioTek, Winooski, VT, USA).




Fig. 1. Cell identification and cell viability assay. (A) α -actinin staining for cell identification. (B) CCK-8 expression levels for the evaluation of cell viability under different cells intervention. * $P < 0.05$, compared with the NG group; # $P < 0.05$, compared with the HG group.

Determination of Cells Apoptosis by Flow Cytometry

Cells apoptosis was detected by double-stained with annexin V-fluorescein isothiocyanate (FITC)/propidium iodide (PI) (BD Pharmingen™, USA), using flow cytometry as described previously (25). In brief, cells in different groups were washed with PBS and detached with trypsin, then re-suspended in binding buffer (1×10^5 cells/ml). Annexin V-FITC (5 μ l) and 5 μ l PI were added and the solution was incubated for 15 min in dark at room temperature. The percentages of cells apoptosis were analyzed using flow cytometry (LSRFortessa, BD Bioscience, San Jose, CA, USA).

Measurement of Intracellular ROS

ROS production was measured using an intracellular ROS assay kit (Biyuntian company, China) and fluorescence microscopy. The cells in different groups were seeded in 24-well plates at 2.0×10^5 /well, and cultured for 24 h. 2',7'-dichlorodihydrofluorescein diacetate (DCHF-DA) was added to each well, which can be oxidized by intracellular ROS. DCHF-DA stock solution (20 mM) was diluted with the medium to a final concentration of 0.1 mM. The cells were incubated at 37°C for 1 h, and washed three times with PBS. Fluorescent compounds, corresponding to intracellular ROS levels, were detected by a fluorescence microscope (ZEISS, Observer A1, Germany) with excitation and emission of 488 nm and 530 nm, respectively. Results were shown by three independent experiments.

Western Blot Analysis

Western blot analyses were conducted to assess the expression levels of the oxidative stress-associated proteins heme oxygenase-1 (HO-1), nuclear

factor E2-related factor 2 (Nrf-2), and the apoptosis-associated proteins B-cell lymphoma 2 (Bcl-2)/Bax, cleaved caspase-3, p-AKT and AKT. A phosphoinositide 3-kinase (PI3K) pathway inhibitor, LY294002 (LY) (5 μ M) (Sigma), was added to the cells incubated with high-glucose and exendin-4, as the HG+Ex+LY Group, for 24 h. In brief, total protein was quantified using a BCA protein assay kit (cwBioTech 02912E, Beijing, PRC) as instructed after lysis of the cardiomyocytes. Equal amounts of proteins (12 μ g) were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). After electrophoresis, proteins were transferred to a polyvinylidene fluoride membrane and blocked with Tris buffered saline containing 5% skim milk powder and 0.05% Tween 20. The corresponding primary antibodies, HO-1 (1:1000, Abcam, Cambridge, MA, USA), Nrf-2 (1:1000, CST, Danvers, MA, USA), Bcl-2/Bax (1:1000, CST) cleaved caspase-3 (1:1000, CST) and p-AKT/AKT (1:1000, CST), were added in series. The membranes were incubated with the diluted antibody preparations overnight at 4°C. After washing, the membranes were incubated with horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (H+L) and goat anti-mouse IgG (H+L) antibodies ($1:10^4$; Jackson, West Grove, PA, USA) for 40 min at room temperature. The blots were visualized using an enhanced chemilumin-essence detection kit (Millipore, Boston, MA, USA). Target proteins were quantified and normalized relative to β -actin (1:1000, Zhongshan, Beijing, PRC).

Statistical Analysis

Data was analyzed by one-way analysis of variance (ANOVA), expressed as mean \pm standard deviation (SD). When a significant difference was identified by ANOVA, a *post-hoc* analysis was performed using the Student-Newman-Keuls test. Data




Fig. 2. Flow cytometry evaluation of cells apoptosis in different groups. (A) NG group. (B) HG group. (C) HG+Ex group. * $P < 0.05$, compared with NG group; # $P < 0.05$, compared with HG group.

were determined using SPSS 13.0 software, and a P -value < 0.05 was considered significant.

Results

Cell Identification and Viability

Observed under an inverted microscope, the cardiomyocytes grew adhering to the wall and showed fusiform or polygon shape, with extension of pseudo-podia and spontaneous beating. α -Actinin staining showed that the cardiomyocyte cell purity reached 90-95% (Fig. 1A). Compared with the NG medium incubation, CCK-8 assay showed that HG incubation reduced cells viability significantly (0.254 ± 0.009 vs. 0.276 ± 0.015 , $P = 0.021$). When exendin-4 was added to HG medium, cells viability improved. Compared to HG medium alone, co-incubation with exendin-4 at 20 nM or 30 nM significantly ameliorated cells viability (0.254 ± 0.009 , $P = 0.010$ vs. 0.278 ± 0.017 and 0.275 ± 0.015 , $P = 0.044$, respectively) (Fig. 1B). Incubation with 20 nM exendin-4 for 24 h significantly improved HG-induced cell injury. Hence, 20 nM exendin-4 was used in further studies.

Effects of Exendin-4 on Cells Apoptosis Analyzed by Flow Cytometry

Flow cytometry analysis demonstrated the effect of exendin-4 on the inhibition of cardiomyocyte apoptosis incubated in HG medium (Fig. 2). Compared with the NG Group, cardiomyocyte apoptosis increased significantly in the HG Group ($P = 0.00$) (Fig. 2D). Although the cell apoptosis rate in the HG+Ex Group was still higher than that in the NG Group, co-incubation with exendin-4 significantly improved HG-induced cell apoptosis in the HG+Ex Group ($P = 0.004$) (Fig. 2D).

Effects of Exendin-4 on ROS Production

Intracellular ROS production was stained with DCHF-DA and determined by fluorescence microscopy. A small amount of ROS was detected in the NG Group (Figs. 3, A and D). Incubated in HG medium, the intracellular ROS level was significantly higher in the HG Group than that in the NG Group ($P = 0.00$) (Fig. 3, B and D). When co-incubated with HG and exendin-4 (HG+Ex Group), the ROS level showed




Fig. 3. Intracellular ROS production assay in different groups. (A) NG group. (B) HG group. (C) HG+Ex group. * $P < 0.05$, compared with the NG group; # $P < 0.05$, compared with the HG group.




Fig. 4. Western blot analysis of expression of oxidative stress-associated proteins. Expression of HO-1 (A) and Nrf-2 protein (B) in different groups. * $P < 0.05$, compared with the NG group; # $P < 0.05$, compared with the HG group.

a significantly decrease (Fig. 3, C and D) compared with the level in the HG Group ($P = 0.00$), although it was still higher than that in the NG Group ($P = 0.002$).

Expression Levels of HO-1 and Nrf-2

HO-1 is a rate-limiting enzyme for oxidative degradation of cellular heme, and also a potent antioxidant enzyme. Nrf-2 is a regulator of numerous genes, encoding antioxidant and cytoprotective factors, such as HO-1. After intervention of HG for 24 h, HO-1 and Nrf-2 protein expression levels were decreased in the HG Group (Fig. 4), as compared with the NG group ($P <$

0.05). But when co-incubated with HG and exendin-4, expression levels of both proteins increased significantly in the HG+Ex Group, when compared with those in the HG Group ($P < 0.05$) (Fig. 4).

Expression Levels of Apoptosis-Associated Proteins

In the HG Group, increased levels of both Bax and cleaved caspase-3 proteins were detected, and Bcl-2 protein expression showed a marked decrease, as compared with the expression levels in the NG Group ($P < 0.05$) (Fig. 5, B, C and D). When co-incubated with exendin-4 in HG medium, Bax and cleaved caspase-3




Fig. 5. Western blot analysis of expression of cell apoptosis-associated proteins. Expression of p-AKT/AKT (A), cleaved caspase-3 (B), Bax (C), and Bcl-2 (D) in different groups. * $P < 0.05$, compared with the NG group; # $P < 0.05$, compared with the HG group; ▲ $P < 0.05$, compared with the HG+Ex group.

significantly declined in expression, and Bcl-2 protein expression displayed a marked elevation, compared with the expression levels in the HG Group ($P < 0.05$). Treatment with LY294002, a PI3K pathway inhibitor, weakened the effects of exendin-4 on the expression of apoptosis-associated proteins in the HG+Ex Group. Bax and cleaved caspase-3 protein levels increased again, and Bcl-2 expression decreased significantly, as compared with the levels in the HG+Ex Group ($P < 0.05$) (Fig. 5, B, C and D). Moreover, p-AKT/AKT level declined significantly in the HG Group ($P < 0.05$), as compared with that in the NG Group. After intervention with exendin-4, p-AKT/AKT level increased in the HG+Ex Group, but the effect was attenuated by LY294002 co-incubation. Compared with the HG+Ex Group, p-AKT/AKT level displayed a significant reduction again in the HG+Ex+LY Group (Fig. 5A).

Discussion

Exendin-4, which has a 53% homology with GLP-1, has become one of the potent anti-diabetic drugs in recent

years. Increasing current interests have focused on its effects on cytoprotection, including the protection of cardiomyocyte (1, 3). As shown by Wen *et al.* (25), exendin-4 can inhibit cardiomyocyte injury in HG incubation, which is achieved through significant inhibition in the expression of receptor for advanced glycation end products (RAGE). In the present study, it was observed that exendin-4 alleviated HG-induced cardiomyocyte apoptosis and improved cells viability. The mechanisms may involve the suppression of oxidative stress *via* the HO-1/Nrf-2 system, and *via* the intervention of apoptosis-associated signal pathways, such as the PI3K/AKT pathway. Firstly, it was shown in this study that exendin-4 ameliorated cardiomyocyte viability and suppressed the HG-induced cell apoptosis. The reduced cell viability by HG incubation was significantly restored in the presence of exendin-4 (Fig. 1B), and there were no obvious differences in cardiomyocyte viability among the NG group and the Ex (20 nM) and Ex (30 nM) incubation groups. In HG+Ex group, cell apoptosis rate and expressions of

pro-apoptosis proteins, Bax and cleaved caspase-3, declined, and the anti-apoptotic protein, Bcl-2, increased in expression level.

Hyperglycemia can activate oxidative stress and causes overproduction of intracellular ROS, which plays a key role in the development and progression of diabetic cardiomyopathy (7). The current study revealed that treatment with exendin-4 decreased HG-induced excessive ROS production, which meant the inhibition of oxidative stress. The result demonstrated the antioxidant properties of exendin-4 in HG environment.

The HO-1/Nrf-2 system has anti-oxidation and anti-apoptosis effects (6, 13). It has also been reported that exendin-4 may interfere with the expressions of HO-1/Nrf-2 and exerts its biological effects in different cell lines (14, 24). In the present study, western blot analysis showed that HO-1 and Nrf-2 protein expression levels of cardiomyocytes treated with exendin-4 were decreased when incubated in high-glucose medium, while the protein levels were elevated significantly in HG+Ex group. HO-1 is the rate-limiting enzyme in heme catabolism. Activation of HO-1 can mediate complex biological functions, exert cytoprotective, anti-inflammatory, anti-oxidative and anti-apoptotic effects (6, 13). Previous studies have revealed the relationship between modulation of HO-1 and the production of ROS in cardiomyocyte and other cell lines (8, 18). Therefore, upregulation of HO-1 *via* exendin-4 might be partly responsible for the anti-oxidative effect, and consequently attenuation of the production of intracellular ROS in the HG+Ex group. Nrf-2 belongs to the Cap'n' Collar (CNC) family of basic leucine zipper (bZip) transcription factors. Once activated, Nrf-2 could be translocated to the nucleus, where it binds to the anti-oxidant response element to up-regulate the expression of numerous cytoprotective phase II detoxifying enzymes and antioxidant genes, such as HO-1 (21). Hence, Nrf-2 is a master transcription factor contributing to HO-1 expression. In the study, exendin-4 exposure significantly elevated the expression of Nrf-2 when compared to HG incubation alone. These results support that exendin-4 may have an effect on the HO-1/Nrf-2 system, and HO-1/Nrf-2 system may play a role in the inhibition of oxidative stress by exendin-4.

The PI3K/AKT signal pathway is central to many cellular survival mechanisms, and some of the biological effects of the HO-1/Nrf-2 system are related to the PI3K/AKT signal pathway (27, 28). Previous studies have indicated that PI3K is the most important regulatory factor for AKT in its upstream pathway, and some phytochemicals could protect against oxidative stress-induced cells damage *via* the PI3K/AKT signal pathway (19, 20). In the present study, we showed that exposure to exendin-4 significantly promoted phosphorylation levels of AKT, which demonstrated AKT activation

and consequent inhibition in the expression of downstream apoptosis-associated genes. Interestingly, LY294002 incubation counteracted the effects of exendin-4 on the AKT pathway and anti-apoptosis in the HG+Ex+LY group. LY294002, a specific inhibitor of PI3K pathway, attenuated the protective effects of exendin-4 in the study, which suggested that exendin-4 might play some biological roles *via* PI3K and its downstream pathways. Taken together, the data suggest that the PI3K/AKT pathway, at least in part, contributed to exendin-4 inhibition of cardiomyocyte apoptosis induced by hyperglycemia.

Based on the findings of this study, it may be speculated that exendin-4 may be a promising therapeutic agent not only for the glucose control, but also for the intervention of diabetic cardiomyopathy.

Acknowledgments

The study was conducted with the support of Capital Medical University, College of Basic Medicine, and the Basic and Clinical Medicine Funding (No. 13-JL58) from Capital Medical University (2013).

References

1. Anagnostis, P., Athyros, V.G., Adamidou, F., Panagioutou, A., Kita, M., Karagiannis, A. and Mikhailidis, D.P. Glucagon-like peptide-1-based therapies and cardiovascular disease: looking beyond glycaemic control. *Diabetes Obes. Metab.* 13: 302-312, 2011.
2. Baggio, L.L. and Drucker, D.J. Biology of incretins: GLP-1 and GIP. *Gastroenterology* 132: 2131-2157, 2007.
3. Chang, G., Zhang, D., Yu, H., Zhang, P., Wang, Y., Zheng, A. and Qin, S. Cardioprotective effects of exenatide against oxidative stress-induced injury. *Int. J. Mol. Med.* 32: 1011-1020, 2013.
4. Drucker, D.J. and Nauck, M.A. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* 368: 1696-1705, 2006.
5. Harkavyi, A. and Whitton, P.S. Glucagon-like peptide 1 receptor stimulation as a means of neuroprotection. *Brit. J. Pharmacol.* 159: 495-501, 2010.
6. Issan, Y., Kornowski, R., Aravot, D., Shainberg, A., Laniado-Schwartzman, M., Sodhi, K., Abraham, N.G. and Hochhauser, E. Heme oxygenase-1 induction improves cardiac function following myocardial ischemia by reducing oxidative stress. *PLoS One* 9: e92246, 2014.
7. Kayama, Y., Raaz, U., Jagger, A., Adam, M., Schellinger, I.N., Sakamoto, M., Suzuki, H., Toyama, K., Spin, J.M. and Tsao, P.S. Diabetic cardiovascular disease induced by oxidative stress. *Int. J. Mol. Sci.* 16: 25234-25263, 2015.
8. Li, B., Kim, D.S., Yadav, R.K., Kim, H.R. and Chae, H.J. Sulforaphane prevents doxorubicin-induced oxidative stress and cell death in rat H9c2 cells. *Int. J. Mol. Med.* 36: 53-64, 2015.
9. Li, H.H., Willis, M.S., Lockyer, P., Miller, N., McDonough, H., Glass, D.J., Patterson, C., Glass, D.J. and Patterson, C. Atrogin-1 inhibits Akt-dependent cardiac hypertrophy in mice *via* ubiquitin-dependent coactivation of Forkhead proteins. *J. Clin. Invest.* 117: 3211-3223, 2007.
10. Lovshin, J.A. and Drucker D.J. Incretin-based therapies for type 2 diabetes mellitus. *Nat. Rev. Endocrinol.* 5: 262-269, 2009.
11. Mangmool, S., Hemplueksa, P., Parichatikanond, W. and Chat-

- tipakorn, N. Epac is required for GLP-1R-mediated inhibition of oxidative stress and apoptosis in cardiomyocytes. *Mol. Endocrinol.* 29: 583-596, 2015.
12. Mannucci, E. and Rotella, C.M. Future perspectives on glucagon-like peptide-1, diabetes and cardiovascular risk. *Nutr. Metab. Cardiovasc. Dis.* 18: 639-645, 2008.
 13. Nepal, S., Kim, M.J., Subedi, A., Lee, E.S., Yong, C.S., Kim, J.A., Kang, W., Kwak, M.K., Arya, D.S. and Park, P.H. Globular adiponectin inhibits ethanol-induced apoptosis in HepG2 cells through heme oxygenase-1 induction. *Biochem. Pharmacol.* 84: 974-983, 2012.
 14. Oeseburg, H., de Boer, R.A., Buikema, H., van der Harst, P., van Gilst, W.H. and Silljé, H.H. Glucagon-like peptide 1 prevents reactive oxygen species-induced endothelial cell senescence through the activation of protein kinase A. *Arterioscler. Thromb. Vasc. Biol.* 30: 1407-1414, 2010.
 15. Robinson, E., Cassidy, R.S., Tate, M., Zhao, Y., Lockhart, S., Calderwood, D., Church, R., McGahon, M.K., Brazil, D.P., McDermott, B.J., Green, B.D. and Grieve, D.J. Exendin-4 protects against post-myocardial infarction remodelling via specific actions on inflammation and the extracellular matrix. *Basic Res. Cardiol.* 110: 20, 2015.
 16. Roul, D. and Recchia, F.A. Metabolic alterations induce oxidative stress in diabetic and failing hearts: different pathways, same outcome. *Antioxid. Redox Signal.* 22: 1502-1514, 2015.
 17. Rubler, S., Dlugash, J., Yuceoglu, Y.Z., Kumral, T., Branwood, A.W. and Grishman, A. New type of cardiomyopathy associated with diabetic glomerulosclerosis. *Am. J. Cardiol.* 30: 595-602, 1972.
 18. Song, B., Zhang, H., Jiang, L., Chi, Y., Tian, J., Du, W.Y.B. and Han, Z. Down-regulation of lipocalin 2 suppresses the growth of human lung adenocarcinoma through oxidative stress involving Nrf2/HO-1 signaling. *Acta Biochim. Biophys. Sin.* 47: 805-814, 2015.
 19. Su, J.D., Yen, J.H., Li, S., Weng, C.Y., Lin, M.H., Ho, C.T. and Wu, M.J. 3',4'-didemethylnobiletin induces phase II detoxification gene expression and modulates PI3K/Akt signaling in PC12 cells. *Free Radic. Biol. Med.* 52: 126-141, 2012.
 20. Tsai, C.Y., Wang, C.C., Lai, T.Y., Tsu, H.N., Wang, C.H., Liang, H.Y. and Kuo, W.W. Antioxidant effects of diallyl trisulfide on high glucose-induced apoptosis are mediated by the PI3K/Akt-dependent activation of Nrf2 in cardiomyocytes. *Int. J. Cardiol.* 168: 1286-1297, 2013.
 21. Tsai, H.Y., Huang, P.H., Lin, F.Y., Chen, J.S., Lin, S.J. and Chen, J.W. Ginkgo biloba extract reduces high-glucose-induced endothelial reactive oxygen species generation and cell adhesion molecule expression by enhancing HO-1 expression via Akt/eNOS and p38 MAP kinase pathways. *Eur. J. Pharm. Sci.* 48: 803-811, 2013.
 22. Tsutsumi, Y.M., Tsutsumi, R., Hamaguchi, E., Sakai, Y., Kasai, A., Ishikawa, Y., Yokoyama, U. and Tanaka, K. Exendin-4 ameliorates cardiac ischemia/reperfusion injury via caveolae and caveolins-3. *Cardiovasc. Diabetol.* 13: 132, 2014.
 23. Wang, C., Chen, X., Ding, X., He, Y., Gu, C. and Zhou, L. Exendin-4 promotes beta cell proliferation via PI3k/Akt signalling pathway. *Cell. Physiol. Biochem.* 35: 2223-2232, 2015.
 24. Yang, H., Li, H., Wang, Z., Shi, Y., Jiang, G. and Zeng, F. Exendin-4 ameliorates renal ischemia-reperfusion injury in the rat. *J. Surg. Res.* 185: 825-832, 2013.
 25. Yi, B., Hu, X., Wen, Z., Zhang, T. and Cai, Y. Exendin-4, a glucagon-like peptide-1 receptor agonist, inhibits hyperglycemia-induced apoptosis in myocytes by suppressing receptor for advanced glycation end products expression. *Exp. Ther. Med.* 8: 1185-1190, 2014.
 26. Younce, C.W., Niu, J., Ayala, J., Burmeister, M.A., Smith, L.H., Kolattukudy, P. and Ayala, J.E. Exendin-4 improves cardiac function in mice overexpressing monocyte chemoattractant protein-1 in cardiomyocytes. *J. Mol. Cell. Cardiol.* 76: 172-176, 2014.
 27. Zhang, G., Wang, Q., Zhou, Q., Wang, R., Xu, M., Wang, H., Wang, L., Wilcox, C.S., Liu, R. and Lai, E.Y. Protective effect of tempol on acute kidney injury through PI3K/Akt/Nrf2 signaling pathway. *Kidney Blood Press Res.* 41: 129-138, 2016.
 28. Zhao, R., Feng, J. and He, G. Hypoxia increases Nrf2-induced HO-1 expression via the PI3K/Akt pathway. *Front. Biosci. (Landmark Ed.)* 21: 385-396, 2016.
 29. Zhao, S.M., Wang, Y.L., Guo, C.Y., Chen, J.L. and Wu, Y.Q. Progressive decay of Ca²⁺ homeostasis in the development of diabetic cardiomyopathy. *Cardiovasc. Diabetol.* 13: 75, 2014.